

# The importance of dogs in eco-epidemiology of Lyme borreliosis: a review

M. BHIDE<sup>1</sup>, M. TRAVNICEK<sup>1</sup>, J. CURLIK<sup>1</sup>, A. STEFANCIKOVA<sup>2</sup>

<sup>1</sup>Department of Epizootology and Infectious Diseases, University of Veterinary Medicine, Kosice, Slovak Republic

<sup>2</sup>Institute of Parasitology, Slovak Academy of Sciences, Kosice, Slovak Republic

**ABSTRACT:** *Borrelia burgdorferi* sensu lato, is endemic in most of the regions in Europe. Pathogen circulates in nature involving ticks vector (mainly *Ixodes ricinus* in Europe) and wide spectrum of reservoir animals like rodents, game animals, birds as well as pets. Considering the close association of a dog and humans, and their similar activities in nature, it is necessary to evaluate the significance of a dog as an important animal in ecology and epidemiology of Lyme borreliosis. Antibody profile in Lyme disease is the most characteristic feature in dogs that helps to evaluate the changes in disease prevalence in particular area and helps to assess the risk factors for human. The article reviews overall eco-epidemiological importance of dogs in Lyme disease surveillance.

**Keywords:** *Borrelia burgdorferi*; dogs; ecology; epidemiology

Lyme disease is the most common arthropod-borne disease of humans in Europe and North America. In Europe, disease has been reported in variety of animal species closely associated with human (Kasbohrer and Schonberg, 1990; Parker and White, 1992; Blowey *et al.*, 1994; Ciceroni *et al.*, 1997; Magnarelli *et al.*, 1997; Stefancikova *et al.*, 2000). Approximately 40 species of mammals and birds have been recognized as a reservoir for *Borrelia burgdorferi* (Bb) (Gern *et al.*, 1998). Among the pet animals dog has been identified as the competent reservoir for *B. burgdorferi* sensu stricto (Mather *et al.*, 1994). Moreover, various researchers have proposed the dog as 'sentinel animal' for the detection of emerging risk areas of Lyme disease (Lindenmayer *et al.*, 1991; Falco *et al.*, 1993; Merino *et al.*, 2000; Bhide *et al.*, 2002). In Europe considerable study on canine Lyme borreliosis has been done with respect to symptoms and seroprevalence. The most com-

mon symptom in dog is migratory arthritis without divergent radiographic findings (Magnarelli *et al.*, 1987). Intermittent lameness can also be seen with several episodes. Other clinical signs consist of anorexia and general malaise. There are some reports of heart block (Levy and Dury, 1988), neurological sign like seizures (Azuma *et al.*, 1993), and fatal kidney failure (Dambach *et al.*, 1997). Although the various symptoms have been reported so far, the diagnosis of Lyme borreliosis in dogs is much more difficult. Recent serological techniques have made the diagnosis easier and more confirmative.

Antibody profile in dogs in Lyme disease is an interesting criterion that can be used for risk assessment of Lyme disease in humans. Purpose of this review is to focus on the possible use of dog as a marker for identification of new developing Lyme disease foci and to elaborate their eco-epidemiological importance.

## Canine Lyme disease serodiagnosis and antibody profile

Till today sensitive serological techniques like ELISA, western blot, immuno fluorescent assay (IFA) etc. are developed to detect and confirm the *B. burgdorferi* infection in dogs. With the help of standard ELISA or IFA, antibody titers can be detected between 4–6 weeks after exposure to infected ticks. Serological tests are also developed to distinguish between early and late stages of *Borrelia* infection. Though the sensitive ELISAs are used widely in humans and pet animals, their specificity is still doubtful. False positive results can be obtained because of vaccination especially in the dogs. In efforts to improve the specificity of serological tests as well as to distinguish between antibodies after *Borrelia* infection and vaccination several strategies have been attempted, for example use of flagellin-enriched (Coleman and Benach, 1987) and purified (Hansen *et al.*, 1988) antigen preparations or specifically VIsE antigen (Liang *et al.*, 2000), and the use of recombinant antigens of *B. burgdorferi* expressed in *Escherichia coli* (Zumstein *et al.*, 1992). Recently, the more specific and confirmative borreliacidal antibody test has been used in dogs (Callister *et al.*, 2000). *B. burgdorferi* infection in humans and other animals results in production of killing (borreliacidal) antibodies. These antibodies are directed against several *B. burgdorferi* proteins including outer surface protein A (OspA), OspB, OspC, decorin binding protein A (DbpA) and outer membrane protein p66 (Scriba *et al.*, 1993; Probert and Lefebvre, 1994; Rousselle *et al.*, 1998; Exner *et al.*, 2000). Borreliacidal antibodies can be detected in dogs one week after attachment of infected tick (Callister *et al.*, 2000). These borreliacidal antibodies not only increase the specificity but also effectively distinguish between early and late Lyme disease.

In untreated infected animal, antibody level increases (IgG), reaching maximum at approximately 90–120 days after tick exposure, and then remain at its level up to one and a half year in the absence of re-exposure (Straubinger, 2000). On the other hand shorter span of anti-*Borrelia* antibodies has been reported in dogs. Moreover, Hovius *et al.* (1999) and Goossens *et al.* (2001) have reported obligatory yearly reinfection to maintain seropositivity in dogs. On the contrary the period of seropositivity in humans after an infection with *B. burgdorferi* is much longer.

## Advantages of canine seroprevalence over serosurvey in other animals

Prediction of potential area for Lyme disease is a difficult task. An epidemiologist may experience more complexities in declaring any new geographical area as emerging risk zone for Lyme borreliosis. The complexity heightens particularly when human case prevalence is low. To identify the endemic area it is also very necessary to study vector-host relationship, vector population and anti-*Borrelia* antibody prevalence in the reservoir hosts. Some researchers have suggested a close association between population/distribution of *Ixodid* ticks and Lyme disease prevalence in humans and dogs (Lissman *et al.*, 1984; Magnarelli *et al.*, 1987). Canine anti-*Borrelia* serosurvey offers a promising tool for targeting areas presenting potential human risk (Rand *et al.*, 1991, 1996). Advantages of canine serosurvey over other animals are: simplicity in sample collection, effective follow up and/or feed back, known history of treatment and vaccination, and greater correlation for Lyme disease risk assessment to human being. On the other hand, tick vector distribution surveys, flagging, small mammal trapping or examination of deer and other wild free-living animals are laborious and time consuming (Eng *et al.*, 1988). Moreover, the seroprevalence in wild animals can not be applied directly to assess the Lyme disease risk to common people who have rare or no contact to forested areas. Measuring the tick density and prevalence of infected ticks in and around cities especially parks, playgrounds and recreational places near human habitat is one of the imperative approach to assess a Lyme risk for common people. As dogs have free and frequent access to such areas, combination of tick density and tick infectivity study with canine seroprevalence can be effective tool to judge the actual Lyme disease risk in the area under study.

## Facts of canine seroprevalence

Till to date considerable work has been done in the field of canine seroprevalence. Anti-*Borrelia* antibodies in dogs have been reported in most of the major European countries (Table 1). Particularly in Slovakia, seropositivity in hunting dogs was 40% whereas; in service and pet dogs positivity observed was 11.80% and 29.40%, respectively (Stefancikova *et al.*, 1996). Difference in seropositivity according to

Table 1. Prevalence of anti-*Borrelia* antibodies in dogs from different parts of the world

Country	County	Method of detection	Prevalence (%)	No. of sample (n)	Reference
Bolivia	Cordillera	ELISA	0.0	43	Ciceroni <i>et al.</i> , 1997
Brazil	Cotia	ELISA	9.7	237	Joppert <i>et al.</i> , 2001
Croatia	Gorski Kotar	ELISA	40.0	10	Poljak <i>et al.</i> , 2000
Czech Republic	Prague	IHA	53.7	169	Sykora <i>et al.</i> , 1990
Germany	–	ELISA	7.2	665	Wittenbrink <i>et al.</i> , 1996
Germany	Berlin	ELISA	10.1	189	Kasbohrer and Schonberg, 1990
Germany	Berlin	IFA	5.8	189	Kasbohrer and Schonberg, 1990
Germany	Bavaria	IFA	35.5	130	Weber <i>et al.</i> , 1991
Israel	–	WB	10.0	40	Beneth <i>et al.</i> , 1998
Italy	Tyrrhenian coast	IFA	0.0	23	Mannelli <i>et al.</i> , 1999
Japan	Tokyo	ELISA	27.3	387	Arashima, 1991
Mexico	Monterrey	IFA	16.0	850	Salinas-Melendez <i>et al.</i> , 1999
Netherlands	(hunting dogs)	ELISA	18.0	448	Goossens <i>et al.</i> , 2001
	(Pet dogs)	ELISA	17.0	75	Goossens <i>et al.</i> , 2001
Slovakia	Kosice	ELISA	26.9	78	Stefancikova <i>et al.</i> , 1996
	Kosice (hunting dogs)	ELISA	45.3	75	Stefancikova <i>et al.</i> , 1998
	(service dogs)	ELISA	18.3	60	Stefancikova <i>et al.</i> , 1998
	(pet dogs)	ELISA	17.6	68	Stefancikova <i>et al.</i> , 1998
Spain	Castilla y Leon	IFA	21.0	308	Delgado and Carmenes, 1995
Spain	Soria	IFA	11.6	146	Merino <i>et al.</i> , 2000
Spain	Leon	IFA	2.10	95	Rojo Vazquez, 1997
Sweden	–	ELISA	3.9	588	Egenvall <i>et al.</i> , 2000
USA	Rhode Island	ELISA	52.0	227	Hinrichsen <i>et al.</i> , 2001
USA	Illinois	ELISA	56.9	1 077	Guerra <i>et al.</i> , 2000
USA	Fort Detrick	ELISA	20.0	440	Sheets <i>et al.</i> , 2000
USA	California	ELISA	2.3	917	Olson <i>et al.</i> , 2000
USA	Alabama	IFA	1.70	579	Wright <i>et al.</i> , 1997
USA	New York	ELISA	49.2	1 446	Falco <i>et al.</i> , 1993
USA	Oklahoma	ELISA	11.7	223	Mukolwe <i>et al.</i> , 1992
USA	Columbia	ELISA	24.3	37	Stockham <i>et al.</i> , 1992
USA	Maine	ELISA	4.34	828	Rand <i>et al.</i> , 1991
USA	Texas	IFA	5.5	2 409	Cohen <i>et al.</i> , 1990
USA	Oklahoma	IFA	18.0	259	Rodgers <i>et al.</i> , 1989
USA	Connecticut	IFA	66.5	155	Magnarelli <i>et al.</i> , 1987
USA	Hudson Valley	IFA	76.3	114	Magnarelli <i>et al.</i> , 1987
USA	New Jersey	IFA	34.7	423	Schulze <i>et al.</i> , 1987
USA	Wisconsin	IFA	53.0	380	Burgess, 1986

use and nature of dogs is also reported by Cohen *et al.* (1990), Stefancikova *et al.* (1998) and Merino *et al.* (2000) whereas, antibody prevalence was not associated with sex and season (Delgado and Carmenes, 1995). Outdoor activity is the prime factor, which

governs percent seropositivity against Lyme borreliosis in any given species of animal. In short, the difference in seroprevalence may due to differences in tasks performed by dogs and therefore the different tick exposition (Daniels *et al.*, 1993). Age de-

pendent variation in seroprevalence of Lyme disease in dogs is reported in Slovakia (Stefancikova *et al.*, 1996), Spain (Merino *et al.*, 2000) and North America (Cohen *et al.*, 1990; Lindenmayer *et al.*, 1991). Some researchers have tried to correlate the seropositivity and geno-phenotypic characteristics of dogs. In dogs with hard type of hairs greater seropositivity against *B. burgdorferi* was reported in comparison to others (Merino *et al.*, 2000). No correlation between size of dog and positivity was reported. Similarly gender is a factor, which does not affect the seropositivity (Magnarelli *et al.*, 1987; Delgado and Carmenes, 1995; Merino *et al.*, 2000). Apart from above explained factors, environment can also play an important role. Dogs living at higher altitude expressed minor seroprevalence in comparison with dogs living in lower region (Lindenmayer *et al.*, 1991). Study in Soria province in Spain by Merino *et al.* (2000) confirmed this hypothesis by comparing the seroprevalences in dogs from other altitudes. All environmental factors ultimately control the tick population in specific area. Tick population governs vector-host relationship as well as tick attachment rate and thus affect the seropositivity in dogs. In the canine surveillance system for Lyme borreliosis in Wisconsin and Illinois (Guerra *et al.*, 2001), seroprevalence pattern by county (0–40%) was significantly correlated with human incidence of Lyme disease and with abundance of tick vector, *Ixodes scapularis*. In the same study a geographic information system (GIS) was used to integrate environmental data with the location of the residences of the dogs to determine environmental risk factors. In Europe environmental risk factors for Lyme disease have been determined using satellite, climatological, and ecological data (Estrada-Pena, 1997; Daniel *et al.*, 1998; Randolph, 2000). Thus, seropositivity in dogs is positively associated with increased tick exposure, time spent outdoor, living in deciduous forested areas etc. Because of close similarity between Lyme disease risk factors of dogs and humans, canine surveillance system is useful method for assessing the risk as well as geographic distribution of Lyme disease.

### **Complement resistance of *Borrelia burgdorferi* and reservoir competence of dog**

Complement-mediated killing of *B. burgdorferi* in hosts have ecological implications as it can determine the reservoir competence (Kurtenbach *et al.*,

1998b; Hovius *et al.*, 2000). The pattern of serum complement sensitivity of different *Borrelia* genospecies matches the known reservoir status of many vertebrate species (Kurtenbach *et al.*, 1998b). Studies indicate that *B. garinii* and *B. valaisiana* are mainly transmitted to ticks by avian hosts whereas, *B. afzelii* is transmitted to tick by rodents (Humair *et al.*, 1995; Kurtenbach, 1998a). *In vitro* canine complement sensitivity test (Hovius *et al.*, 2000) against three different *Borrelia* strains (B31, *B. burgdorferi* sensu stricto; pKo, *B. afzelii*; and A87S, *B. garinii*) showed B31 and pKo as resistant species to dog complement than A87S. It was observed that *Borrelia* isolates differ in their ability to activate complement and resist killing by serum bactericidal activity (Brade *et al.*, 1992). Though there is no extensive study available to compare species specific complement sensitivity of *Borrelia* and reservoir competence of dog, one can extrapolate the available complement sensitivity results to propose reservoir status of dog for particular *Borrelia* species (Hovius *et al.*, 2000). Such a correlation was made previously in rodents and squirrels by Kurtenbach *et al.* (1998b). Rodent complement resistance of *B. afzelii* parallels the prime transmission competence of rodent species (Humair *et al.*, 1995) and squirrels (Craine *et al.*, 1997). Furthermore, complement mediated lyses of *B. garinii* explains why the European rodents are insufficient reservoir for European *B. garinii* strains, while its resistance to pheasant complement makes clear the concept of reservoir competence ability of pheasant for the same *Borrelia* genospecies. In case of an incompetent reservoir sika deer active killing of *Borrelia* by complement takes place (Nelson *et al.*, 2000). Similarly lysis of *Borrelia* regardless of genospecies correlates the incompetent reservoir nature of deer explained by Jaenson and Talleklint (1992).

### **Lyme disease risk assessment for pet owners and hunters**

Overall Lyme disease risk assessment data compiled in various reviews and reports (Flisiak and Zabicka, 1995; Arteaga and Garcia-Monco, 1999; Werner *et al.*, 2001), indicate the morbidity exceeds 100 cases per 100 000 inhabitants per year in central Europe. Comparatively higher prevalence observed in outdoor workers than indoor workers in southwest Sweden (Werner *et al.*, 2001) and Spain (Arteaga and Garcia-Monco, 1999) indicates posi-

tive correlation between human contact with tick vector and Lyme disease risk. Only measurement of human contact with a tick is not sufficient criteria to assess the risk of Lyme disease. Apart from this criterion, the population of reservoir competent domestic, wild as well as pet animals in particular area is necessary to study.

There are many controversies about zoonotic importance of pet animals as far as Lyme disease is concern. Even if some authors have had put forth hypothesis about greater risk of Lyme disease to pet owners (Mather *et al.*, 1994), there is no concrete evidence of direct infection from pet animals or dogs to human. Survey in the Netherlands by Goossens *et al.* (2001), showed no correlation between sole ownership of dogs and seropositivity against Lyme disease. However, recently a case of one and a half year old girl suffering from gonarthitis has been reported by Zajadacz and Juskiewicz (2002) with high antibody titre. The girl was never in the forest and had no contact with animals except a pet dog. Authors suggested the most possible transmission of Lyme disease from pets to the girl.

Hunting dogs usually carry infected ticks from the forest. Loosely attached ticks as well as infected females from dog drop near human habitat. Female ticks lay eggs in the spring, which hatches to larvae. These Ixodid larvae preferentially feed on small mammals and rodents. Presence of rodent population in and around human habitat facilitates feeding of larvae and nymphs and consequently helps in establishment of the tick population. Rodents are known competent reservoir from which Ixodid larvae acquire *Borrelia* infection. In the following spring larvae moult into nymphs, with an acquired infection from rodents. Ixodid nymphs have wide host range including dogs and humans. Nymphs moult to adult in fall and act as the most important source of infection for dogs. Transmission of *Borrelia* from Ixodid ticks to dogs, cats and human has been reported (Smith *et al.*, 1993). Higher seroprevalence (33%) in the domestic cats does not exclude the importance of this pet animal in Lyme disease epidemiology (Magnarelli *et al.*, 1990).

To conclude, screenings of dog for seropositivity is good indicator of actual and present risk of Lyme disease in particular area due to shorter span of anti-*Borrelia* IgG antibodies. Dogs stay seropositive for a much shorter period after an infection with *Borrelia*. On the other hand the seropositivity in other animals as well as in humans persists for several years. Similar seroprevalence in hunting

dogs and humans particularly in hunters (Goossens *et al.*, 2001) illuminates close relation and linked epidemiological aspects of Lyme disease. Evolution and establishment of Lyme disease focus may occur quickly due to favourable climatic conditions and geoeological suitability of central Europe for tick vectors. Hunting dogs can serve as seroindicators and/or sentinel for identifying new focuses as well as assessing the changes in endemicity of well known focuses of Lyme disease.

## Acknowledgements

Special thanks are due to Dr. S. B. Barbuddhe, Sr. Scientist, ICAR Goa, India for his constructive comments and correction of the manuscript. We thank Ing. A. Eibenová for reviewing the English of the manuscript.

## REFERENCES

- Arashima Y. (1991): Anti-*Borrelia burgdorferi* antibody in dogs: Lyme disease as zoonosis. *Rinsho Byori*, 39, 869–874.
- Arteaga Perez F., Gracia-Monco Carra J.C. (1999): Risk factors associated with the presence of antibodies against *Borrelia burgdorferi*. *Rec. Clin. Esp.*, 199, 136–141.
- Azuma Y., Kawamura K., Isogai H., Isogai E. (1993): Neurological abnormalities in two dogs suspected Lyme disease. *Microbiol. Immunol.*, 37, 325–329.
- Baneth G., Breitschwerdt E.B., Hegarty B.C., Pappalardo B., Ryan J. (1998): A survey of tick-borne bacteria and protozoa in naturally exposed dogs from Israel. *Vet. Parasitol.*, 74, 133–142.
- Bhide M.R., Curlik J., Travnicek M., Stefancikova A. (2002): Hunting Dogs, Sentinel of Lyme Disease. In: *Compendium, International Conference on Actual Problems in Zoonoses*, September 2002, Brno, Czech Republic.
- Blowey R.W., Carter S.D., White A.G., Barnes A. (1994): *Borrelia burgdorferi* infection in U.K. cattle: a possible association with digital dermatitis. *Vet. Rec.*, 135, 577–578.
- Brade V., Kleber I., Acker G. (1992): Differences of two *Borrelia burgdorferi* strains in complement activation and serum resistance. *Immunobiology*, 185, 453–465.
- Burgess E.C. (1986): Natural exposure of Wisconsin dogs to the Lyme disease spirochete (*Borrelia burgdorferi*). *Lab. Anim. Sci.*, 36, 288–290.

- Callister S.M., Jobe D.A., Schell R.F., Lovrich S.D., Onheiber K.L., Korshus J.B. (2000): Detection of borrelia-cidal antibodies in dogs after challenge with *Borrelia burgdorferi* infected *Ixodes scapularis* ticks. *J. Clin. Microbiol.*, *38*, 3670–3674.
- Ciceroni L., Bartoloni A., Ciarrochi S., Pinto A., Paradisi F. (1997): Serological survey for antibodies to *B. burgdorferi* in sheep, goats and dogs in cordillera province. Bolivia *J. Vet. Med. B*, *44*, 133–137.
- Cohen N.D., Carter C.N., Thomas M.A., Angulo A.B., Bugster A.K. (1990): Clinical and epidemiologic characteristics of dogs seropositive for *Borrelia burgdorferi* in Texas: 110 cases (1988). *J. Am. Vet. Med. Assoc.*, *197*, 893–898.
- Coleman J.L., Benach J.L. (1987): Isolation of antigenic components from the Lyme disease spirochete: their role in early diagnosis. *J. Infect. Dis.*, *155*, 756–765.
- Craine N.G., Nuttall P.A., Marriott A.C., Randolph S.E. (1997): Role of gray squirrels and pheasants in the transmission of *Borrelia burgdorferi* sensu lato, the Lyme disease spirochete, in the UK. *Folia Parasitol.*, *44*, 155–160.
- Dambach D.M., Smith C.A., Lewis R.M., Van Winkle T.J. (1997): Morphologic, immunohistochemical, and ultrastructural characterization of a distinctive renal lesion in dogs putatively associated with *Borrelia burgdorferi* infection: 49 cases (1987–1992). *Vet. Pathol.*, *34*, 85–96.
- Daniel M., Kolar J., Zeman P., Pavelka K., Sadlo J. (1998): Predictive map of *Ixodes ricinus* high-incidence habitats and a tick-borne encephalitis risk assessment using satellite data. *Exp. Appl. Acarol.*, *22*, 417–433.
- Daniels T.J., Fish D., Levine J.F. (1993): Canine exposure to *Borrelia burgdorferi* and prevalence of *Ixodes dammini* (Acari: Ixodidae) on deer as a measure of Lyme disease risk in the northeastern United States. *J. Med. Entomol.*, *30*, 171–178.
- Delgado S., Carmenes P. (1995): Seroepidemiological survey for *Borrelia burgdorferi* (Lyme Disease) in dogs from northwestern of Spain. *Eur. J. Epidemiol.*, *11*, 321–324.
- Egenvall A., Bonnett B.N., Gunnarsson A., Hedhammar A., Shoukri M., Bornstein S., Artursson K. (2000): Seroprevalence of granulocytic *Ehrlichia* spp. and *Borrelia burgdorferi* sensu lato in Swedish dogs 1991–94. *Scand. J. Infect. Dis.*, *32*, 19–25.
- Eng T.R., Wilson M.L., Spielman A., Lastavica C.C. (1988): Greater risk of *Borrelia burgdorferi* infection in dogs than in people. *J. Infect. Dis.*, *158*, 1410–1411.
- Estrada-Pena A. (1997): Epidemiological surveillance of tick populations: A model to predict the colonization success of *Ixodes ricinus* (Acari: Ixodidae). *Eur. J. Epidemiol.*, *13*, 573–580.
- Exner M.M., Wu X., Blanco D.R., Miller J.N., Lovett M.A. (2000): Protection elicited by native outer membrane protein Oms66 (p66) against host adapted *Borrelia burgdorferi*: conformational nature of bactericidal epitopes. *Infect. Immun.*, *68*, 2647–2654.
- Falco R.C., Smith H.A., Fish D. (1993): The distribution of canine exposure to *Borrelia burgdorferi* in a Lyme disease endemic area. *Am. J. Public Health*, *83*, 1305–1310.
- Flisiak R., Zabicka J. (1995): Epidemiological situation of Lyme borreliosis in Europe. *Przegl. Epidemiol.*, *49*, 375–379.
- Gern L., Estrada-Pena A., Frandsen J., Gray T., Jaenson F., Jongejan O., Khal E., Korenberg R., Mehl R., Nuttall P. (1998): European reservoir hosts of *Borrelia burgdorferi* sensu lato. *Zentbl. Bakteriol. Parasitenkd. Infektkrankh. Hyg. Abt. 1 Orig.*, *287*, 196–204.
- Goossens H.A.T., van dan Bogaard A.E., Nohlmans M.K.E. (2001): Dogs as sentinel for human Lyme borreliosis in The Netherlands. *J. Clin. Microbiol.*, *39*, 844–848.
- Guerra M.A., Walker E.D., Kitron U. (2000): Quantitative approach for the serodiagnosis of canine Lyme disease by the immunoblot procedure. *J. Clin. Microbiol.*, *38*, 2628.
- Guerra M.A., Walker E.D., Kitron U. (2001): Canine surveillance system for Lyme borreliosis in Wisconsin and northern Illinois: geographic distribution and risk factor analysis. *Am. J. Trop. Med. Hyg.*, *65*, 546–552.
- Hansen K., Hindersson P., Pedersen N.S. (1988): Measurement of antibodies to the *Borrelia burgdorferi* flagellum improves serodiagnosis in Lyme disease. *J. Clin. Microbiol.*, *26*, 338–346.
- Hinrichsen V.L., Whitworth U.G., Breitschwerdt E.B., Hegarty B.C., Mather T.N. (2001): Assessing the association between the geographic distribution of deer ticks and seropositivity rates to various tick-transmitted disease organisms in dogs. *J. Am. Vet. Med. Assoc.*, *218*, 1092–1097.
- Hovius J.W.R., Hovius K.E., Oei A., Houwers D.J., van Dam A.P. (2000): Antibodies against specific proteins of and immobilizing activity against three strains of *Borrelia burgdorferi* sensu lato can be found in symptomatic but not in infected asymptomatic dogs. *J. Clin. Microbiol.*, *38*, 2611–2621.
- Hovius K.E., Rijpekma S.G., Westers P., van der Zeijst B.A.M., van Asten F.J.A.M., Houwers D.J. (1999): A serological study of cohorts of young dogs, naturally exposed to *Ixodes ricinus* ticks, indicates seasonal reinfection by *Borrelia burgdorferi* sensu lato. *Vet. Q.*, *21*, 16–20.
- Humair P.F., Peter O., Wallich R., Gern L. (1995): Strain variation of Lyme disease spirochetes isolated from

- Ixodes ricinus* ticks and rodents collected in two endemic areas in Switzerland. J. Med. Entomol., 32, 433–438.
- Jaenson T.G.T., Talleklint L. (1992): Incompetence of roe deer as reservoir of the Lyme borreliosis spirochete. J. Med. Entomol., 29, 813–817.
- Joppert A.M., Hagiwara M.K., Yoshinari N.H. (2001): *Borrelia burgdorferi* antibodies in dogs from Cotia county, Sao Paulo State, Brazil. Rev. Inst. Med. Trop. Sao Paulo, 43, 251–255.
- Kasbohrer A., Schonberg A. (1990): Serologic studies of the occurrence of *Borrelia burgdorferi* in domestic animals in Berlin (West). Berl. Munch. Tierarztl. Wochenschr., 103, 374–378.
- Kurtenbach K., Peachey M., Rijpkema S.G.T., Hoodless A.N., Nuttall P.A., Randolph S.E. (1998a): Differential transmission of the genospecies of *Borrelia burgdorferi* sensu lato by game birds and small rodents in England. Appl. Environ. Microbiol., 64, 1169–1174.
- Kurtenbach K., Sewell H.S., Ogden N.H., Randolph S.E., Nuttall P.A. (1998b): Serum complement sensitivity as a key factor in Lyme disease ecology. Infect. Immunol., 66, 1248–1251.
- Levy S.A., Dury P.H. (1988): Complete heart block in a dog seropositive for *Borrelia burgdorferi*. J. Vet. Intern. Med., 2, 138–144.
- Liang F.T., Jacobson R.H., Straubinger R.K., Grooters A., Philipp M.T. (2000): Characterization of a *Borrelia burgdorferi* VlsE invariable region useful in canine Lyme disease serodiagnosis by enzyme-linked immunosorbent assay. J. Clin. Microbiol., 38, 4160–4166.
- Lindenmayer J.M., Marshall D., Onderdonk A.B. (1991): Dogs as sentinel for Lyme disease in Massachusetts. Am. L. Public Health, 81, 1448–1455.
- Lissman B.A., Bosler E.M., Camay H., Ormistan B.G., Benach J.L. (1984): Spirochete associated arthritis (Lyme disease) in a dog. J. Am. Vet. Med. Assoc., 185, 219–220.
- Magnarelli L.A., Anderson F.J., Schreier A.B., Ficke C.M. (1987): Clinical and serologic studies of canine borreliosis. J. Am. Vet. Med. Assoc., 191, 1089–1094.
- Magnarelli L.A., Anderson J.F., Levine H.R., Levy S.A. (1990): Tick parasitism and antibodies to *Borrelia burgdorferi* in cats. J. Am. Vet. Med. Assoc., 197, 63–66.
- Magnarelli L.A., Flavell R.A., Padula S.J., Anderson J.F., Fikrig E. (1997): Serologic diagnosis of canine and equine Borreliosis: Use of recombinant antigen in enzyme-linked immunosorbent assay. J. Clin. Microbiol., 35, 169–173.
- Mannelli A., Cerri D., Buffrini L., Rossi S., Rosati S., Arata T., Innocenti M., Grignolo M.C., Bianchi G., Iori A., Tolari F. (1999): Low risk of Lyme borreliosis in a protected area on the Tyrrhenian coast, in central Italy. Eur. J. Epidemiol., 15, 371–377.
- Mather T.N., Fish D., Coughlin R.T. (1994): Competence of dogs as reservoir for Lyme disease spirochetes (*Borrelia burgdorferi*). J. Am. Vet. Med. Assoc., 205, 186–188.
- Merino F.J., Serrano J.L., Saz J.V., Nebreda T., Gegundez M., Beltran M. (2000): Epidemiological characteristics of dogs with Lyme borreliosis in the province of Soria (Spain). Eur. J. Epidemiol., 16, 97–100.
- Mukolwe S.W., Kocan A.A., Wyckoff J.H. 3rd. (1992): Serological survey for Lyme disease in domestic dogs and white-tailed deer from Oklahoma. Ann. N.Y. Acad. Sci., 653, 172–177.
- Nelson D.R., Rooney S., Miller N.J., Mather T.N. (2000): Complement – mediated killing of *Borrelia burgdorferi* by nonimmune sera from sika deer. J. Parasitol., 86, 1232–1238.
- Olson P.E., Kallen A.J., Bjorneby J.M., Creek J.G. (2000): Canines as sentinels for Lyme disease in San Diego County, California. J. Vet. Diagn. Invest., 12, 126–129.
- Parker J.L., White K.K. (1992): Lyme borreliosis in cattle and horses: a review of literature. Cornell Vet., 82, 253–274.
- Poljak I., Troselj-Vukic B., Miletic B., Morovic M., Ruzic-Sabljić E., Vucemilovic A., Materljan E. (2000): Low seroprevalence of Lyme borreliosis in the forested mountainous area of Gorski Kotar, Croatia. Croat. Med. J., 41, 433–436.
- Probert W.S., Lefebvre R.B. (1994): Protection of C3H/HeN mice from challenge with *Borrelia burgdorferi* through active immunization with OspA, OspB, or OspC, but not with OspD or the 83-kilodalton antigens. Infect. Immun., 62, 1920–1926.
- Rand P.W., Smith R.P., Lacombe E.H. (1991): Canine seroprevalence and the distribution of *Ixodes dammini* in an area of emerging Lyme disease. Am. J. Public Health, 81, 1331–1334.
- Rand P.W., Lacombe E.H., Smith R.P., Gensheimer K., Dennis D.T. (1996): Low seroprevalence of human Lyme disease near a focus of high entomologic risk. Am. J. Trop. Med. Hyg., 55, 160–164.
- Randolph S.E. (2000): Tick and tick borne disease system in space and from space. Adv. Parasitol., 47, 217–243.
- Rodgers S.J., Morton R.J., Baldwin C.A. (1989): A serological survey of *Ehrlichia canis*, *Ehrlichia equi*, *Rickettsia rickettsii* and *Borrelia burgdorferi* in dogs in Oklahoma. J. Vet. Diagn. Invest., 1, 154–159.
- Rojo Vazquez J. (1997): Seroprevalence of the infections caused by *Borrelia burgdorferi* and *Rickettsia conorii* in humans and dogs in primary health care of San Andreas del Rabanedo. Rev. Esp. Salud Publica, 71, 173–180.
- Rousselle J.C., Callister S.M., Schell R.F., Lovrich S.D., Jobe D.A., Marks J.A., Wieneke C.A. (1998): Borreliacidal

- antibody production against outer surface protein C of *Borrelia burgdorferi*. *J. Infect. Dis.*, 178, 733–741.
- Salinas-Melendez J.A., Avalos-Ramirez R., Riojas-Valdez V.M., Martinez-Munoz A. (1999): Serological survey of canine borreliosis. *Rev. Latinoam. Microbiol.*, 41, 1–3.
- Schulze T.L., Bosler E.M., Shisler J.K., Ware I.C., Lakat M.F., Parkin W.E. (1987): Prevalence of canine Lyme disease from an endemic area as determined by serosurvey. *Zentralbl. Bakteriol. Mikrobiol. Hyg. A*, 263, 427–434.
- Scriba M., Ebrahim J.S., Schlott T., Eiffert H. (1993): The 39-kilodalton protein of *Borrelia burgdorferi*: a target for borreliacidal human monoclonal antibodies. *Infect. Immun.*, 61, 4523–4526.
- Sheets J.T., Rossi C.A., Kearney B.J., Moore G.E. (2000): Evaluation of a commercial enzyme-linked immunosorbent assay for detection of *Borrelia burgdorferi* exposure in dogs. *J. Am. Vet. Med. Assoc.*, 216, 1418–1422.
- Smith R.P. Jr., Rand P.W., Lacombe E.H., Telford S.R., 3rd., Rich S.M., Piesman J., Spielman A. (1993): Norway rats as reservoir hosts for Lyme disease spirochetes on Monhegan Island, Maine. *J. Infect. Dis.*, 168, 687–691.
- Stefancikova A., Skardova I., Petko B., Janovska D., Cyprichova V. (1996): Antibodies IgG to *Borrelia burgdorferi* in dogs from Kosice region (in Slovak). *Vet. Med. – Czech*, 41, 83–86.
- Stefancikova A., Tresova G., Petko B., Skardova I., Sestakova E. (1998): ELISA comparison of three whole cell antigens of *Borrelia burgdorferi* sensu lato in serological study of dogs from area of Kosice, Eastern Slovakia. *Ann. Agric. Environ. Med.*, 5, 25–30.
- Stefancikova A., Stepanova G., Petko B., Nadzamova D., Sestakova E., Skardova I., Leinstein R. (2000): Prevalence of antibodies to *Borrelia burgdorferi* in horses of east Slovakia. *Vet. Med. – Czech*, 45, 227–231.
- Stockham S.L., Schmidt D.A., Curtis K.S., Schauf B.G., Tyler J.W., Simpson S.T. (1992): Evaluation of granulocytic ehrlichiosis in dogs of Missouri, including serologic status to *Ehrlichia canis*, *Ehrlichia equi* and *Borrelia burgdorferi*. *Am. J. Vet. Res.*, 53, 63–68.
- Straubinger R.K. (2000): Lyme borreliosis in dogs. In: Carmichael L.E. (ed.): *Recent Advances in Canine Infectious Diseases*. IVIS Publication.
- Sykora J., Minar J., Petrikova O., Vokoun P., Gojda M. (1990): The occurrence of antibodies to *Borrelia* in dogs. *Vet. Med. (Praha)*, 35, 251–256.
- Weber A., Heim U., Schafer R. (1991): Incidence of antibodies to *Borrelia burgdorferi* in dogs in small animal practice in North Bavaria. *Berl. Munch. Tierarztl. Wochenschr.*, 104, 384–386.
- Werner M., Nordin P., Arnholm B., Elgefors B., Krantz I. (2001): *Borrelia burgdorferi* antibodies in outdoor and indoor workers in south-west Sweden. *Scand. J. Infect. Dis.*, 33, 128–131.
- Wittenbrink M.M., Failing K., Krauss H. (1996): Enzyme-linked immunosorbent assay and immunoblot analysis for detection of antibodies to *Borrelia burgdorferi* in dogs. The impact of serum absorption with homologous and heterologous bacteria. *Vet. Microbiol.*, 48, 57–68.
- Wright J.C., Chambers M., Mullen G.R., Swango L.J., D'Andrea G.H., Boyce A.J. (1997): Seroprevalence of *Borrelia burgdorferi* in dogs in Alabama, USA. *Prev. Vet. Med.*, 127–131.
- Zajadacz B., Juszkiewicz A. (2002): Gonarthrosis in the course of Lyme disease in a one and a half year old child. *Wiad. Lek.*, 55, 243–246.
- Zumstein G., Fuchs R., Hofmann A., Preac-Mursic V., Soutschek E., Wilske B. (1992): Genetic polymorphism of the gene encoding the outer surface protein A (OspA) of *Borrelia burgdorferi*. *Med. Microbiol. Immunol.*, 181, 57–70.

Received: 03–04–14

Accepted after corrections: 04–03–05

---

*Corresponding Author*

Dr. Mangesh Bhide, University of Veterinary Medicine, Department of Epizootology and Infectious Diseases, Komenskeho 73, 041 81 Košice, Slovak Republic  
Fax +421 55 632 36 66, e-mail: mangeshbhide@hotmail.com

---